Digital Sequence Information: A Looming Disaster or Hidden Opportunity for Positive Change?



GROW Webinar August 26, 2021












Dr. Amber Hartman Scholz, Deputy to the Director





The Convention on Biological Diversity has 3 objectives




The CBD was not enough....


Realm Living Planet Indices

As seen in the map below (figure 22), populations are in decline in all realms, but declines are especially pronounced in the three tropical realms. Here, average vertebrate abundance in 2014 was less than half of what it was in 1970. The LPI indicates that the Neotropical realm, covering South and Central America, and the Caribbean, has suffered the most dramatic decline at 89% loss compared to 1970. Nearctic and Palearctic populations are faring slightly better with declines of 23% and 31%.

The Palearctic Living Planet Index The average abundance for 2,866 terrestrial and freshwater populations (representing 576 species). These populations declined by 31% (-6% to -50%) between 1970 and 2014'.



PALEARCTIC




The Nearctic Living Planet Index The average abundance for 2,417 terrestrial and

freshwater populations (representing 875 species). These populations declined by 23% (+11% to -47%) between 1970 and 2014'.

NEOTROPICAL

Figure 22: The Living Planet Index for each realm The white line in the LPI graph shows the index values and the shaded areas represent the statistical certainty surrounding the trend (95%). Threat data is available for 3,789 – just under a quarter – of the populations in the global LPI'. This is explored in chapter 2.




NEARCTIC

The Neotropical Living Planet Index The average abundance for 1,040 populations (representing 689 species) in the Neotropical biogeographic realm. Vertebrate populations declined by an average of 4.8% annually between 1970 and 2014, translating to an overall decline of 89%. This represents the most dramatic change in any biogeographic realm'.

Tuger (1900 = 1) 190 2000 2010

The Afrotropical Living Planet Index
The average abundance for 1,115 terrestrial and
freshwater Afrotropical populations (representing
320 species). These populations declined by
56% between 1970 and 2014. The decrease was
particularly steep in the 1980s'.



The Indo-Pacific Living Planet Index The average abundance for 1,083 terrestrial and freshwater populations (representing 488 species) in the Indo-Pacific biogeographic realm. The Indo-Pacific Index experienced the second most pronounced decline, after the Neotropical index: 64% overall between 1970 and 2014.

COP 10 October 2010, Nagoya, Japan

A horse trade / Kuhhandel: The Nagoya Protocol & The Aichi Targets



Aichi Targets



Understand values



Reduce pollution



Enhance resilience



Mainstream biodiversity



Reduce invasive spp.



Implement Nagoya Prot.



Address incentives



Minimize reef loss



Revise NBSAPs



Sustainable production



Protected areas



Respect and conserve TK



Halve rate of loss



Prevent extinctions



Improve knowledge



Sustainable fisheries



Conserve gene pool



Mobilize resources



Manage within limits



Restore ecosystems

The Nagoya Protocol

Nagoya Protocol on Access to Genetic Resources and the Fair and Equitable Sharing of Benefits Arising from their Utilization (NP) to the Convention on Biological Diversity (CBD)

Aims

- to provide **legal certainty** for the actors involved in the use and international exchange of genetic resources & traditional knowledge
- to specify **means for sharing benefits** gained from the use of genetic materials & traditional knowledge


Need to know


<u>Genetic resources</u> - plants, animal, fungi, microbes

- biochemical compounds


- DNA/RNA

Use = non-commercial and commercial






Procedure to comply with CBD and NP





Plant scientists have a complicated ABS situation



Aubry, S- Front. Plant Sci., 30 August 2019 https://doi.org/10.3389/fpls.2019.01046



COP 15 October 2021 / May 2022, Kunming, China

Another horse trade?
Global Biodiversity Framework &
Digital Sequence Information



The Aichi Targets expired in 2020

Up next:

- **2020 Global Biodiversity Framework**
- > set regional goals, be more specific!



TABLE 2 Trends shown by agreed indicators of progress towards the 2010 biodiversity target

	Z	Trends in extent of selected biomes, ecosystems, and habitats	Most habitats in most parts of the world are declining in extent, although forest area expands in some regions, and the loss of mangroves has slowed significantly, except in Asia.
	Z	Trends in abundance and distribution of selected species	Most species with limited population size and distribution are being further reduced, while some common and invasive species become more common. 禁業業 (but limited number of taxa assessed)
	K	Change in status of threatened species	The risk of extinction increases for many threatened species, although some species recovery programmes have been very successful.
	Z	Trends in genetic diversity of domesticated animals, cultivated plants, and fish species of major socio-economic importance	It is likely that the genetic variety of cultivated species is declining, but the extent of such decline and its overall impacts are not well understood. *** (although many case studies with a high degree of certainty are available)
		Coverage of protected areas	There has been a significant increase in coverage of protected areas, both terrestrial and marine, over the past

Ecosystem integrity and ecosystem goods and services

Status and trends of the components of biological diversity

4	Marine Trophic Index	Despite intense pressure the Marine Trophic Index has shown a modest increase globally since 1970. However there is substantial regional variation with declines being recorded in half of the marine areas with data. Although the global increases may indicate a recovery it is more likely a consequence of fishing fleets expanding their areas of activity, thus encountering fish stocks in which larger predators have not yet been removed in large numbers.
Z	Connectivity – fragmentation of ecosystems	Most terrestrial and aquatic ecosystems are becoming increasingly fragmented, despite an increased recognition of the value of corridors and connections, especially in climate change adaptation.
	Water quality of aquatic	Most parts of the world are likely to be suffering from declines in water quality, although quality in some

areas has improved through control of point-source pollution

7	Nitrogen deposition	Human activity has doubled the rate of creation of reactive nitrogen on the planet's surface. Pressure on biodiversity from nutrient pollution continues to increase, although some measures to use nutrients more efficiently, to reduce their release into water and the atmosphere, are beginning to show positive effects.
Commence of the	war and the terror to a	The control of the form of the control of the contr

7	Trends in invasive alien species	The number and rate of spread of allen species is increasing in all continents and all ecosystem types. ************************************

Sustainable use

2	Area of forest, agricultural and aquaculture ecosystems under sustainable management	There are considerable efforts under way to increase the extent of areas of land under sustainable management. Regional efforts on sustainable forest management are expected to contribute to this. Traditional agricultural practices are being maintained and revitalized as the demand for ethical and healthy products increases. However, these are still relatively small niches and major efforts are required to substantially increase the areas under sustainable management.
7	Ecological footprint and related concepts	The ecological footprint of humanity is increasing. Efforts at increasing resource efficiency are more than compensated by increased consumption by a growing and more prosperous human population.


Status of traditional knowledge, innovations and practices

A =	Status and trends of linguistic	A large number of minority languages are believed in danger of disappearing, and linguistic diversity is very
	diversity and numbers of speakers	likely declining.
	of indigenous languages	(although case studies with a high degree of certainty are available)


Status of access and benefit sharing


benefit-sharing to be Working Group on Access and Benefit-sharing.	7 bene		The need and possible options for additional indicators are being examined by the Ad Hoc Open-ende Working Group on Access and Benefit-sharing.	d
--	--------	--	---	---

Official development assistance (ODA) provided in support of the	The volume of ODA for biodiversity has increased over the past few years.
Convention	常常常










decade. However, many ecological regions, particularly in marine ecosystems, remain underprotected, and

the management effectiveness of protected areas remains variable.

DSI under the CBD: what is happening during the 2018–2020 intersessional period?




Rohden and Scholz. PLANTS, PEOPLE, PLANET, First published: 05 May 2021, DOI: (10.1002/ppp3.10198)

Nature crisis: Talks resume on global plan to protect biodiversity

By Matt McGrath Environment correspondent









Preparations for the Post-2020 Biodiversity Framework

Third meeting of the Open-ended Working Group on the Post-2020 Global Biodiversity Framework

UN negotiators have resume

Join the WG2020 3 meeting

nature and species for the next uecaue.


The draft Global Biodiversity Framework aims to conserve at least 30% of the world's land and oceans.



iren GmbH

Why is there a call for benefit sharing from DSI? There has been an explosion in sequencing since the Nagoya Protocol


Growth of DNA Sequencing





Leibniz-Institut • DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH

The biotechnological tools have also changed significantly since 2010! Synthetic biology & Gene editing



**

**

**

**

**

Biological features Genomic simplicity

Physiological simplicity

Homologous recombination

Gene expression strength
Absence of transgene silencing

Photoautotrophic growth

Polycistronic gene expression

Biological transgene containment

Available tools and techniques

Ease of genetic manipulation Speed of genetic manipulation

Genetic part characterization

Genetic & metabolic network models

Genetic part assembly

Genetic part insulation

Gene expression control

Demonstrated applications

Synthetic subcompartments

©2019 by American Society of Plant Biologists

Synthetic genetic circuits

Synthetic genomes

Synthetic metabolic pathways

Genetic parts



CRISPR-Cas9 "genetic scissors" 2020 Nobel Prize in Medicine

https://hub.jhu.edu/2021/01/29/crispr-cas9-gene-dimmer/

Christian R. Boehm and Ralph Bock Plant Physiol. 2018;179:794-802

But the data ecosystem is NOT like the Nagoya Protocol


- 212 billion sequences (from every country in the world)
- Entire DSI dataset downloaded 34 million times a year
- 10-15 million users (in every country in the world)
- Promulgated via 2,000 downstream databases
- Half a million publications and patent applications

Most DSI does NOT come from low-/middle-income countries

What is the country of origin for non-human NSD?


1.	China	18.23 %
2.	United States	17.39 %
3.	Canada	9.10 %
4.	Japan	7.24 %
5.	India	3.46 %
6.	Australia	2.66 %
7.	Mexico	2.54 %
8.	Brazil	2.30 %
9.	Germany	1.83 %
10.	Spain	1.58 %

52% of DSI comes from 4 countries



There are 10-15 million users of the core DSI databases (INSDC). They live in every country in the world.





INSDC infrastructure costs: \$50 million per year → \$3-5 per user ~50% of users live in countries that do not contribute to infrastructure costs.


Their use is currently "subsidized".



Which countries use more than they provide?

How does database usage compare to provided sequences?

1.	Lebanon	27.68
2.	Ukraine	23.84
3.	Belarus	22.88
4.	Iraq	21.41
5.	Colombia	19.79
6.	Algeria	19.54
7.	Pakistan	19.37
8.	Republic of Korea	18.81
9.	Belgium	17.83
10.	Poland	17.47



Technical observations

- 16% of DSI entries reported country information
- 44% of DSI entries did not report a country but should have
- 4% of NSD entries linked to GR

Genebanks are well-positioned to improve DSI metadata!!!



PubMed ID

Original GR



What about the commercial use of DSI?

- 20% of DSI in INSDC comes from patent-associated NSD directly deposited by a patent office
- 2. NSD itself is more frequently listed as reference material than the basis for the invention
- 3. No country information associated with DSI in databases even if listed in patent application
- 4. Patent NSD often receives a new AN even if it originated in INSDC

Fall 2019:


There is a DSI deal that will be made someday soon.

How can the scientific community contribute?





WiLDSI project: asking scientists how to maintain open access and engage with benefit-sharing



Epidemiology




Wildlife conservation




Reducing waste in pig feed fish



PANGAEA: Data Publisher for Earth & Environmental Science



Insect taxonomy & **Loss of pollinators**




Antimicrobial resistance



Tropical ecosystem research & climate change adaptation



Poaching & Wildlife trade



Mollusk barcoding



Fungal diversity & nutrient cycling



Viral metagenomics & bioprospecting



Freshwater Biodiversity and Evolution






5 requirements for successful science


- 1. Open Access
- 2. Simplicity: Learn from the Nagoya Protocol
- 3. "Future-Proof"
- 4. Legal Certainty
- 5. Opt-in Genetic Resources



Not all DSI is under the CBD... "Universal solution"?



DSMZ









https://www.youtube.com/watch?v=rO_pAQDDHII

Scholz, A.H., Hillebrand, U., Freitag, J., Cancio, I., dos S. Ribeiro, C., Haringhuizen, G., Oldham, P., Saxena, D., Seitz, C., Thiele, T., van Zimmeren, E. (2020) Finding Compromise on ABS & DSI in the CBD: Requirements & Policy Ideas from a Scientific Perspective. doi: 10.13140/RG.2.2.35180.80001.



5 Open-Access Policy Options for DSI

Option 1 : Micro-Levy (multilateral)

Option 2: Membership Fee (multilateral)


Option 3: Cloud-based fees (multilateral)

Option 4: Commons Licenses for DSI (bilateral with multilateral licenses)

Option 5: Blockchain Metadata, open DSI (bilateral with multilateral elements)



March 2021: DSI policy options synthesized by the CBD Secretariat



So which DSI option is the best?

Let's learn more from data on DSI provision and use

http://wildsi.ipk-gatersleben.de/

Scholz et al. Myth-busting the provider-user relationship for digital sequence information. BioRxiv pre-print. DOI: 10.1101/2021.08.02.454535.



Using the scientific I providing and using

AF307324; SV 1; linear; genomic RNA; STD; VRL; 643 BP.

25-OCT-2000 (Rel. 65, Created) 25-OCT-2000 (Rel. 65, Last updated, Version 1)

Castelo dos Sonhos virus small RNA segment, nucleocapsid protein gene, partial cds.

Novel St Associat DOI; 10.100 Infecti

Castelo dos Sonhos virus

Viruses; Riboviria; Negarnaviricota; Polyploviricotina; Ellioviricetes; Bunyavirales; Hantaviridae; Mammantavirinae; Orthohantavirus.

Rollin P.E

This example \rightarrow Johnson A.: Providing country: Brazil "Genetic in Using country: Peru

.CO:2-Y.

Ksiazek T.G.,

Cristhopher D. Cruz, Brett M. Fors [2] Carolina Guevara, V. Albe

Author affiliations: US Naval Medical Resear Peru (C.D. Cruz, B.M. Forshey, D.L. Blazes, Laguna-Torres, E.S. Halsey); Servicio Departa Europepmc; PMC136083; 11907216. Cochabamba, Bolivia (E. Vallejo, R. Agudo); C Enfermedades Tropicales, Santa Cruz, Bolivia (Naval Medical Research Center, Silver Spring, N Kochel)

```
DSMZ
            Leibniz-Institut • DSMZ-Deutsch
```

```
Johnson A.M., de Souza L.T.M., Ferreira I.B., Pereira L.E., Ksiazek T.G.,
Rollin P.E., Peters C.J., Nichol S.T.;
Submitted (22-SEP-2000) to the INSDC.
Special Pathogens Branch/DVRD/NCID, Centers for Disease Control and
Prevention, 1600 Clifton Road, Atlanta, GA 30333, USA
```

```
MD5; bbb10a3ac321b8c3a4de9dd395da5290.
EuropePMC; PMC3381544; 21801642.
EuropePMC; PMC3503106; 23055565.
EuropePMC; PMC3598921; 23259834.
EuropePMC; PMC3901500; 24447689.
```

/1 \6/2

```
Location/Qualifiers
                1..643
source
                /organism="Castelo dos Sonhos virus"
                /chromosome="S seament"
                /mol type="genomic RNA"
                /country="Brazil: castelo dos Sonhos, Mato Grasso State"
                 note="detected in human serum sample"
                /db xref="taxon:139033"
```

EF534082), Neembucu virus (S: DQ345763), Alta Paraguay virus (S: DQ345762), Itapua virus (S: DQ345765), Araraquara virus (ARAV; S: AF307325, M: AF307327), Araucaria virus strain HPR/03-99 (S: AY740630), Jabora virus strain Akp8048 (S: JN232080), Juquitiba virus strain Olfo 777 (S: GU213198), and Castelo dos Sonhos virus (CASV; S: AF307324, M: AF307326).

Sequences were aligned by using ClustalW (www. clustal.org) with manual adjustments, and pairwise genetic distances were calculated by using MEGA4.0 (26). For phylogenetic analysis, maximum-likelihood (ML) and Bayesian approaches were used. ML phylogenetic rees were estimated by using PhyML (27,28) and 100 bootstrap replications to place confidence intervals at HPS in Brazil"; nodes. Phylogenetic reconstructions were also conducted in MrBayes version 3.1 (29,30) under the general time reversible + Γ + proportion invariant model of evolution, with 1 million Markov Chain Monte Carlo generations, and sampling every 100 generations with a burn-in of 25,000. Puumula virus S and M segments were included as outgroups in the phylogenetic reconstructions.


Results

Patient Identification

During January 2008–June 2009, serum samples from 372 febrile patients reporting to clinics in Chapare Province, Bolivia (Figure 1) were tested for serologic evidence of recent infection by ≥ 1 hantaviruses. Of these 372 patients, 199 (53.5%) were male patients with a median age of 31 years (range 7-95 years). IgM against ANDV (n = 8) or LNV (n = 1) antigen was identified in acute-phase or convalescent-phase samples from 9 (2.4%) patients. No evidence of recent arbovirus infection was detected in these samples. Of the 9 patients with IgM against hantaviruses, 7 (77.8%) were male patients with a median age of 32 years range 15-49 years). Three of the 9 patients were positive for hantavirus RNA.


Patient 1 (FVB0554) was an 18-year-old man (student)

79 countries are using Kenyan DSI






Kenyan scientists are using DSI from 83 countries






DSI from Mexico is used by 75 countries






Scientists from Mexico are using DSI from 83 countries



Eritrea




94 countries are using Indian DSI






Indian scientists are using DSI from 150 countries






57 countries are using Philippine DSI






Philippine scientists are using DSI from 32 countries






Countries are both providers AND users of DSI – often roughly equally



DSI is used more by "locals" than by "foreigners"




DSI Scientific Network

- Founded in November 2020, 36 members, 16 Countries
- Members are DSI providers, users, researchers
- Mission: to contribute to policymakers and other stakeholders' understanding of DSI, its applications and contributions to research supporting biodiversity conservation and public health, as well as the global benefits of open access to DSI databases.
- **Goal:** Members recognise that finding a solution, which balances the value of **open access** for researchers with the importance of **benefit sharing**, including monetary benefits, is essential to deliver a global biodiversity framework that meets the priorities of different stakeholders and contributes to the Sustainable Development Goals.

A de-coupled multilateral framework for DSI-BS

- 1. A bilateral or "tagged" mechanism will disincentivize use of LMIC-sourced DSI
- 2. Access should be de-coupled from benefitsharing (facilitate rather than compete)
- 3. Virtuous cycle: Recognize and incentivize national contributions (esp. from LMICs)
- 4. Core focus: Conservation, sustainable use and bioeconomic innovation (complement the GBF goals)
- 5. Capacity building for science to improve DSI exploitation (rather than regulatory mechanisms)



Acknowledgments



Mehmood Ghaffar IPK former



Dr. Matthias Lange IPK



Pia Katherina-Habekost IPK



Jens Freitag, IPK



Fabian Rohden, DSMZ former







Dr. Guy Cochrane EMBL-EBI



Dr. Blaise Alako EMBL-EBI



Upneet Hillebrand, DSMZ former



Distr.
GENERAL
CBD/DSI/AHTEG/2020/1/4
31 January 2020
ENGLISH ONLY

CBD

AD HOC TECHNICAL EXPERT GROUP ON DIGITAL SEQUENCE INFORMATION ON GENETIC RESOURCES Montreal, Canada, 17-20 March 2020

COMBINED STUDY ON DIGITAL SEQUENCE INFORMATION IN PUBLIC AND PRIVATE DATABASES AND TRACEABILITY

https://www.cbd.int/doc/c/1f8f/d793/57cb114ca40cb6468f479584/dsi-ahteg-2020-01-04-en.pdf

Thank you!

amber.h.scholz@dsmz.de

New DSI workshop series: http://io.aibs.org/nagoya.










https://www.dsmz.de/collection/nagoya-protocol/digital-sequence-information http://wildsi.ipk-gatersleben.de/





Leibniz-Institut • DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH